




a division of  
Amgrow Australia Pty Ltd

# PRODUCT INFORMATION SHEET

## Auxinone

*Active Constituents: 0.075 g/L Naphthalene Acetic Acid, 0.075 g/L Indole Acetic Acid, 2.25 g/L Vitamin B1 Thiamine*

Auxinone is a blend of root hormone stimulants and vitamins for use in turf, sugarcane and propagation situations. Auxinone is a root growth formula used to increase early root initiation in treated plants. The two auxin types (IAA/NAA) present in Auxinone promotes the development of new root tips first, then followed by early leaf formation. Vitamin B1 is naturally produced in plant leaves and transported to roots where it stimulates root growth and development. Auxinone also contains a high concentration of vitamin B1.



### Benefits:

- Encourages rapid prolific root development
- More uniform emergence after planting
- Vigorous early leaf development

### Pack Size:

Available in 500mL and 20 L packs



Transplants

### Product Characteristics:

Colour	Specific Gravity	pH
Pale Red	0.97	4.7



Cane Set



a division of  
Amgrow Australia Pty Ltd

# Auxinone

## Directions for Application (refer to product label for more detailed instructions)

Auxinone can be applied in a tank mix with agricultural insecticides/fungicides.

CROP	RATE	CRITICAL COMMENTS
Turf (bowling green)	1600 mL in 1600 litres of water	Apply to standard bowling green
Turf (golf green)	100 mL per 100 litres of water/100m <sup>2</sup>	Apply at 3-4 weekly intervals during the growing season.
Turf (sporting field & oval)	10 litres per 1000 litres of water /ha	Apply at 3-4 weekly intervals during the growing season.
Sugar Cane	100 mL per 100 litres of water	Apply thoroughly to the billets prior to planting. Can be applied with fungicide solution
Propagation (cuttings, pipings, transplants)	10 mL per 10 litres of water	Dip or spray planting material with solution before planting.

